

Note to Readers: If you need assistance accessing items in this Supplemental Material, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Table of Contents for Supplemental Material

Maternal Polybrominated Diphenyl Ether (PBDE) Exposure and Thyroid Hormones in Maternal and Cord Sera: The HOME Study, Cincinnati, USA

Ann M. Vuong, Glenys M. Webster, Megan E. Romano, Joseph M. Braun, R. Thomas Zoeller, Andrew N. Hoofnagle, Andreas Sjödin, Kimberly Yolton, Bruce P. Lanphear, and Aimin Chen

Table S1. Comparison of published studies on polybrominated diphenyl ether and thyroid hormone levels in pregnancy.

Table S2. Coefficients of variation for thyroid hormone and antibody assays.

Table S3. Thyroid hormone levels, HOME Study.

Table S4. Spearman rank correlation matrix for polybrominated diphenyl ether congeners and thyroid hormones, HOME Study.

Table S5. Spearman rank correlation matrix for thyroid hormone levels in maternal and cord sera, HOME Study.

Table S6. Adjusted associations between maternal polybrominated diphenyl ether concentrations and maternal and cord sera levels of thyroid antibodies, HOME Study.

Table S7. Associations between polybrominated diphenyl ether concentrations and select thyroid hormones in maternal and cord sera by thyroid antibody levels, HOME Study.

Figure S1. β -coefficients and 95% CIs from regression models for associations of BDE-28 and BDE-47 (ng/g lipid) quartiles and thyroid hormones in cord serum. All models adjusted for maternal age, race/ethnicity, education, parity, family income, smoking status, alcohol consumption, infant gender, mode of delivery, gestational age at serum collection, and total serum PCB concentrations. P trend was obtained by using the median value in each quartile as a continuous variable in the linear regression models.

References